



Medication Guidelines

IMPORTANT! Please carefully review this medication list and **consult with your prescribing provider** before discontinuing any blood thinners/anticoagulants or diabetic medications. **DO NOT** make up or double up on any missed medications after your procedure.

	10 DAYS BEFORE PROCEDURE	5 DAYS BEFORE PROCEDURE	4 DAYS BEFORE PROCEDURE	3 DAYS BEFORE PROCEDURE	2 DAYS BEFORE PROCEDURE	1 DAY BEFORE PROCEDURE	DAY OF PROCEDURE
Blood Thinners/Anticoagulants							
• Ticlid® (ticlopidine)	DO NOT TAKE						
• Coumadin® (warfarin) • Effient® (prasugrel) • Plavix® (clopidogrel)	DO NOT TAKE						
• Brilinta® (ticagrelor)	DO NOT TAKE						
• Aggrenox® • Arixtra® (fondaparinux) • Eliquis® (apixaban) • Persantine® (dipyridamole) • Pletal® (cilostazol) • Savaysa® (edoxaban) • Xarelto® (rivaroxaban)	DO NOT TAKE						
• Fragmin® (dalteparin) • Lovenox® (enoxaparin) • Pradaxa® (dabigatran)	DO NOT TAKE						
Vitamins & Supplements							
• Fish Oil • Iron • Multivitamins • St. John's Wort • Vitamin E • All other supplements	DO NOT TAKE						
Weight Loss (see more under Diabetic & Weight Loss Meds)							
• Phentermine/phendimetrazine (all brands)	DO NOT TAKE						
Pain Relievers							
• Aspirin (ASA) • Tylenol® (acetaminophen)	Continue taking <i>Nothing by mouth 4hrs prior to appointment arrival time</i>						
• NSAIDs (Advil, Aleve, ibuprofen, naproxen, diclofenac, meloxicam)	Continue taking in small doses <i>Nothing by mouth 4hrs prior to appointment arrival time</i>						

	DAY BEFORE PROCEDURE	DAY OF PROCEDURE
Blood Pressure - ACE Inhibitors & ARBs		
<ul style="list-style-type: none"> • Accupril® (quinapril) • Aceon® (perindopril) • Altace® (ramipril) • Atancand® (candesartan) • Avapro® (irbesartan) • Benicar® (olmesartan) • Captopril • Cozaar® (losartan) • Diovan® (valsartan) • Edarbi® (azilsartan) • Eprosartan • Fosinopril • Lotensin® (benazepril) • Mavik® (trandolapril) • Micardis® (telmisartan) • Moexipril • Prinivil® (lisinopril) • Vasotec® (enalapril) • Zestril® (lisinopril) 	Take as usual	DO NOT TAKE
Blood Pressure - Diuretics		
<ul style="list-style-type: none"> • Bumex® (bumetanide) • Chlorthalidone • Demadex® (torsemide) • Diuril® (chlorothiazide) • Edecrin® (ethacrynic acid) • Indapamide • Lasix® (furosemide) • Microzide® (hydrochlorothiazide) • Metolazone 	Take as usual	DO NOT TAKE
Blood Pressure - Beta Blockers & Calcium Channel Blockers		
<ul style="list-style-type: none"> • Adalat® (nifedipine) • Bystolic® (nebivolol) • Calan® (verapamil) • Cardene® (nicardipine) • Cardizem LA® (diltiazem) • Corgard® (nadolol) • Covera-HR® (verapamil) • Dynacirc® (isradipine) • Inderal LA® (propranolol) • Innopran XL® (propranolol) • Lopressor® (metoprolol) • Nimotop® (nimodipine) • Norvasc® (amlodipine) • Plendil® (felodipine) • Procardia® (nifedipine) • Sectral® (acebutolol) • Sular® (nisoldipine) • Tenormin® (atenolol) • Tiazac® (diltiazem) • Toprol XL® (metoprolol) • Verelan PM® (verapamil) • Zebeta® (bisoprolol) 	Take as usual	Take as usual



Medication Guidelines - Diabetic

IMPORTANT: If diabetic, consult with your prescribing provider before discontinuing any diabetic medications

	7 DAYS BEFORE PROCEDURE	1 DAY BEFORE PROCEDURE	DAY OF PROCEDURE
Diabetes and/or Weight Loss Injectable & Oral Medications (GLP-1 Agonists)			
<ul style="list-style-type: none"> • Adlyxin® (lixisenatide) • Bydurean® (exanatide ER) • Byetta® (exenatide) • Mounjaro® (tirzepatide) • Ozempic® (semaglutide) • Rybelsus® (semaglutide) 	<ul style="list-style-type: none"> • Saxenda® (liraglutide) • Symlin® (pramlintide) • Tanzeum® (albiglutide) • Trulicity® (dulaglutide) • Victoza® (liraglutide) • Wegovy® (semaglutide) 	<p>If diabetic, consult with your prescribing provider before discontinuing any diabetic medications</p> <p>GLP-1 medications taken weekly should be stopped 7 days prior to procedure</p>	<p>Follow Clear Liquid Diet 24 hours prior to procedure arrival time</p> <p>NOTHING BY MOUTH 4 hours prior to procedure arrival time.</p> <p>DO NOT TAKE GLP-1 taken daily should be stopped the day of procedure</p> <p>AFTER procedure, if eating, take usual dose if due that day</p>

	96 HOURS BEFORE PROCEDURE	72 HOURS BEFORE PROCEDURE	1 DAY BEFORE PROCEDURE	DAY OF PROCEDURE
Diabetes and/or Heart Failure Oral Medications (SGLT2 Inhibitors)				
• Steglatro® (ertugliflozin)	DO NOT TAKE			DO NOT TAKE AFTER procedure, if eating, take usual dose
• Farxiga® (dapagliflozin) • Invokana® (canagliflozin) • Jardiance® (empagliflozin)	DO NOT TAKE			DO NOT TAKE AFTER procedure, if eating, take usual dose
Other Diabetic Medications				
Metformin & metformin combo meds <ul style="list-style-type: none"> • Actoplus Met® (pioglitazone/metformin) • Avandamet® (metformin) • Glucophage® (metformin) • Glucovance® (glyburide/metformin) • Invokamet® (canagliflozin/metformin) • Janumet® (sitagliptin/metformin) • Jentaduo® (linagliptin/metformin) • Kazano® (alogliptin/metformin) • Kombiglyze® (saxagliptin/metformin) • Metaglip® (glipizide/metformin) • PrandiMet® (metformin) • Repaglin® (repaglinide/metformin) • Riomet® (metformin) • Synjardy® (empagliflozin/metformin) • Xigduo XR® (dapagliflozin/metformin) 			DO NOT TAKE EVENING DOSE	DO NOT TAKE Resume usual dose day after procedure
Non-metformin meds <ul style="list-style-type: none"> • Actos® (pioglitazone) • Avandia® (rosiglitazone) • Amaryl® (glimepiride) • Avandaryl® (glimepiride/rosiglitazone) • DiaBeta® (glyburide) • Duetact® (pioglitazone/glimepiride) • Glucotrol® (glipizide) • Glyset® (miglitol) • Januvia® (sitagliptin) • Onglyza® (saxagliptin) • Prandin® (repaglinide) • Precose® (acarbose) • Starlix® (nateglinide) • Tradjenta® (linagliptin) 			Take as usual	DO NOT TAKE AFTER procedure, if eating, take usual dose

IMPORTANT: Diabetic patients should check blood sugar 4x daily if taking insulin.

	1 DAY BEFORE PROCEDURE			DAY OF PROCEDURE		
	AM dose	PM dose	Bed dose	AM dose	PM dose	Bed dose
Insulin						
<ul style="list-style-type: none"> • Lantus® (insulin glargine) • Levemir® (insulin detemir) • Novolin® (isophane) • Soliqua (lixisenatide/insulin glargine) • Tresiba® (insulin degludec) • Xultophy® (liraglutide/insulin degludec) 			Take half of usual dose if normally taken at bedtime	Take half of usual dose after procedure if normally taken in AM	Take usual dose after procedure if eating	Take usual dose
<ul style="list-style-type: none"> • Apidra® (insulin glulisine) • Humalog®, Humalog S® (insulin lispro) • Humulin® (regular insulin) • Novolin® Mix 70/30 • Novolog® (insulin aspart) • Novolog® Mix 70/30 • Pre-mixed insulin: 75/25 • Regular insulin 	Take half of usual fixed dose OR cover carbs with usual carb rotation		Take half of usual fixed dose	DO NOT TAKE	Take usual dose after procedure if eating & check blood sugars	Take usual dose at bedtime if eating & check blood sugars
• Insulin Pens	Same as above					
• Insulin Pumps	Continue insulin without change. Administer additional insulin per carbohydrate or other protocol.					